

## PET/CT rebatable items

### Lung-Solitary Pulmonary Nodule

Whole body FDG PET study, performed for evaluation of a solitary pulmonary nodule where the lesion is considered unsuitable for transthoracic fine needle aspiration biopsy, or for which an attempt at pathological characterisation has failed. (R)

### NSCLC

Whole body FDG PET study, performed for the staging of proven non-small cell lung cancer, where curative surgery or radiotherapy is planned. (R)

### Brain

FDG PET study of the brain for evaluation of suspected residual or recurrent malignant brain tumour based on anatomical imaging findings, after definitive therapy (or during ongoing chemotherapy) in patients who are considered suitable for further active therapy. (R)

### Colorectal

Whole body FDG PET study, following initial therapy, for the evaluation of suspected residual, metastatic or recurrent colorectal carcinoma in patients considered suitable for active therapy. (R)

### Melanoma

Whole body FDG PET study, following initial therapy, performed for the evaluation of suspected metastatic or recurrent malignant melanoma in patients considered suitable for active therapy. (R)

### Epilepsy

FDG PET study of the brain, performed for the evaluation of refractory epilepsy which is being evaluated for surgery. (R)

### Ovarian

Whole body FDG PET study, following initial therapy, performed for the evaluation of suspected residual, metastatic or recurrent ovarian carcinoma in patients considered suitable for active therapy. (R)

### Uterine/Cervix

Whole body FDG PET study, for the further primary staging of patients with histologically proven carcinoma of the uterine cervix, at FIGO stage IB2 or greater by conventional staging, prior to planned radical radiation therapy or combined modality therapy with curative intent. (R)

### Uterine/Cervix

Whole body FDG PET study, for the further staging of patients with confirmed local recurrence of carcinoma of the uterine cervix considered suitable for salvage pelvic chemoradiotherapy or pelvic exenteration with curative intent. (R)

### Oesophageal/GEJ

Whole body FDG PET study, performed for the staging of proven oesophageal or GEJ carcinoma, in patients considered suitable for active therapy. (R)

### Head/Neck

Whole body FDG PET study performed for the staging of biopsy-proven newly diagnosed or recurrent head and neck cancer. (R)

### Head/Neck (Residual)

Whole body FDG PET study performed for the evaluation of patients with suspected residual head and neck cancer after definitive treatment, and who are suitable for active therapy. (R)

### Metastatic SCC- unknown primary

Whole body FDG PET study performed for the evaluation of metastatic squamous cell carcinoma of unknown primary site involving cervical nodes. (R)

### NHL- Initial staging

Whole body FDG PET study for the initial staging of indolent non-Hodgkin's lymphoma where clinical, pathological and imaging findings indicate that the stage is I or IIA and the proposed management is definitive radiotherapy with curative intent. (R)

### Lymphoma-Initial staging

Whole body FDG PET study for the initial staging of newly diagnosed or previously untreated Hodgkin's or non-Hodgkin's lymphoma (excluding indolent non-Hodgkin's lymphoma). (R)

### Lymphoma Therapy Response

Whole body FDG PET study to assess response to first line therapy either during treatment or within three months of completing definitive first line treatment for Hodgkin's or non-Hodgkin's lymphoma (excluding indolent non-Hodgkin's lymphoma). (R)

### Lymphoma- Restaging

Whole body FDG PET study for restaging following confirmation of recurrence of Hodgkin's or non-Hodgkin's lymphoma (excluding indolent non-Hodgkin's lymphoma). (R)

### Lymphoma- Post chemotherapy

Whole body FDG PET study to assess response to second-line chemotherapy when stem cell transplantation is being considered, for Hodgkin's or non-Hodgkin's lymphoma (excluding indolent non-Hodgkin's lymphoma). (R)

### Bone/Soft Tissue Sarcoma

Whole body FDG PET study for initial staging of patients with biopsy-proven bone or soft tissue sarcoma (excluding gastrointestinal stromal tumour) considered by conventional staging to be potentially curable. (R)

### Residual/Recurrent Sarcoma

Whole body FDG PET study for the evaluation of patients with suspected residual or recurrent sarcoma (excluding gastrointestinal stromal tumour) after the initial course of definitive therapy to determine suitability for subsequent therapy with curative intent. (R)